

COFFEE PLANT WITH REDUCED α -D-GALACTOSIDASE ACTIVITY

Cross-Reference to Related Applications

This application is a continuation of International application PCT/EP02/09148 filed August 15, 2002, the entire content of which is expressly incorporated herein by reference thereto.

Background Art

The present invention relates to the modification of galactomannans present in green coffee beans by reducing the endogenous level of α -D-galactosidase activity. In particular the present invention pertains to a plant cell having a reduced α -D-galactosidase activity and to a plant harboring such a plant cell.

In coffee grains, cell wall polysaccharides account for approximately 48% of mature coffee bean dry weight, and of these, mannans represent approximately half. These polysaccharides are essentially insoluble in purified form and have very low galactose branching (Bradbury and Haliday, J. agric. Food Chem. **38** (1990), 389-392). Mannan polymers are acknowledged to be the main reason for the large losses of original green coffee weight encountered during preparation of soluble coffee drinks. The losses occur either when insoluble material remains as sediments during initial extraction or when precipitates and gels form during storage of coffee liquors. Mannans have also been shown to be the principal component responsible for cloudiness and precipitation during standing of coffee beverages.

In some plants, the degree of galactose branching on the mannan chains has been found to partially depend on the activity of the α -D-galactosidase (EC 3.2.1.22). This enzyme is capable of releasing α -1,6-linked galactose units from galactomannans stored in plant seed storage tissue or maturation (Buckeridge and Dietrich, Plant Sci. **117** (1996), 33-43). In addition, the accumulation of galactomannans having a very low galactose/mannose ratio in some plant endosperms or cotyledon tissues has been shown to correlate to peak α -D-galactosidase activity during maturation of these tissues and to the hardening and drying thereof (Kontos and Spyropoulos, Plant Physiol. Biochem. **34** (1996), 787-793).

α -D-galactosidases activity has also been associated with the capacity to remove galactose residues, i.e., α -1,6-linked to galactomannan polysaccharides, which brings about a decreased solubility of these polymers (McCleary, Carb. Res. **92** (1981), 269-285).

Furthermore, the removal of galactose side chains from galactomannans seems to increase the capacity thereof to interact with other polysaccharides, e.g., xanthans in guar, with a concomitant formation of complex gel. Galactose branching on coffee grain mannans decreases from approximately 40 % in young grains to the low level found in the mature grains during maturation. Concurrently, α -D-galactosidase enzyme activity increases during coffee grain maturation.

The coffee α -D-galactosidase cDNA has been cloned (Zhu and Goldstein, Gene 140 (1994), 227-231). According to the information derived therefrom, the mature coffee bean α -D-galactosidase is presumed to be composed of 363 amino acids and is synthesized as a preproenzyme of 420 residues. Following biosynthesis, two protease cleavages then remove a secretion signal (38 residues) and another signal peptide (19 residues) to produce the protein exhibiting the N-terminal amino acid sequence characteristic of the active enzyme.

In view of the known effects of galactomannans on the preparation and/or storability of soluble coffee, there is a need in the art to improve this situation. More specifically, there is a need to provide an improved method for preparing soluble coffee while concurrently obviating the drawbacks known when storing coffee liquors.

Summary of the Invention

The present invention satisfies the above-mentioned needs by providing a coffee plant cell, and a coffee plant, respectively, wherein the galactose branching in the galacto-mannans is increased. The increased galactose branching is in turn achieved by reducing the endogenous level of α -D-galactosidase activity.

The coffee plant cell generally contains a nucleic acid that is transcribed to a ribonucleic acid which is antisense to mRNA, or a part thereof, derived from the α -D-galactosidase gene of the cell. Advantageously, the nucleic acid is under the control of a constitutive or inducible promoter, such as a coffee *csp1* promoter.

The invention also relates to a coffee plant containing a coffee plant cell as described herein. This plant can produce coffee beans which form yet another embodiment of the invention. These coffee beans can be used in a method for preparing a coffee beverage or a method for increasing the solubility of coffee particles

Detailed Description of the Preferred Embodiments

According to a preferred embodiment of the invention, a coffee plant cell, and a coffee plant, respectively, may be obtained by reducing the endogenous level of α -D-galactosidase activity so that the galactose branching in the galacto-mannans is increased. The reduction of α -D-galactosidase activity may be achieved by conventional methods of mutation and selection using the techniques available in the art. Thus, plant cells may be subjected to mutagenic treatments, such as by exposing them to chemicals or radiation that bring about an alteration of the cell's DNA. The cells thus treated are subsequently screened for the desired property.

According to another preferred embodiment, such a reduced endogenous level of α -D-galactosidase activity is obtained by introducing a construct into a coffee plant cell, containing a nucleic acid that is transcribed into an antisense copy of the mRNA encoded by the α -D-galactosidase gene, or to a part thereof.

To this end, the antisense copy of the mRNA encoded by the α -D-galactosidase gene may be any ribonucleic acid capable of forming dimers under physiological conditions, i.e., to hybridize with the mRNA encoded by the α -D-galactosidase gene under conditions prevailing in the cell. Thus, the antisense copy does not need to be a 100 % homologue to the corresponding counterpart, but rather needs to provide sufficient binding for forming a dimer.

Consequently, antisense copies (and the corresponding nucleic acids from which they are transcribed), that are modified by substitution, deletion and/or insertion of nucleotides are well within the context of the present invention. In this respect, it will also be appreciated that the antisense copy may represent a full counterpart to the mRNA encoded by the α -D-galactosidase gene, that is, it may provide a RNA molecule having essentially the same length as the mRNA encoding the α -D-galactosidase polypeptide. On the other hand the antisense copy may only cover a part of the mRNA encoding the α -D-galactosidase polypeptide.

The nucleic acid encoding a ribonucleic acid, antisense to the mRNA encoded by the α -D-galactosidase gene, or to a part thereof, may be under the control of a constitutive or an inducible promoter, so that the level of the antisense RNA may be conveniently controlled. However, in all cases the level of the antisense copy should be sufficiently high so as to reduce the number of mRNA copies encoding the α -D-galactosidase polypeptide accessible for the ribosomes.

According to a preferred embodiment the promoter utilized is the coffee *csp1* promoter, which gives a sufficiently high transcription rate.

The present invention therefore provides for a modified coffee plant cell and a coffee plant, respectively, wherein the level of α -D-galactosidase activity has been reduced such that eventually the galactose branching on galacto-mannans is increased. According to a preferred embodiment the plant is a transgenic plant, the cells of which harbor a construct capable to provide an antisense copy of the mRNA derived from the α -D-galactosidase gene or a part thereof.

The present invention also provides a method for preparing soluble coffee, which comprises the step of using coffee beans derived from a plant exhibiting a reduced α -D-galactosidase activity. In this respect the present invention also provides a method for increasing the solubility of coffee by increasing the galactose branching.

This invention aims to increase in the solubility of coffee galacto-mannans by increasing their galactose branching. The strategy adopted is to reduce the endogenous level of α -D-galactosidase activity, preferably by introducing an antisense copy of its cDNA under the control of the coffee *csp1* promoter.

This *csp1* promoter has already been characterized (Marraccini *et al.*, Plant Physiol. Biochem. **37** (1999), 273-282) and controls the expression a gene encoding coffee 11S storage protein. A cassette containing said promoter was constructed and introduced into the T-DNA region of a binary vector of transformation, which derives from the pTiT37 plasmid (Bevan, Nucl. Acids Res. **12** (1984), 8711-8721). This recombinant vector was introduced in *Agrobacterium tumefaciens*, which was used to transform coffee explants. Coffee plants harboring the T-DNA inserted in their genome were regenerated and analyzed for the α -D-galactosidase activity in their grains.

I. Analysis of the α -D-galactosidase activity in coffee grains during maturation

Fruits were harvested at different stage of maturation (age is expressed as weeks after flowering: WAF) from *Coffea arabica* variety Caturra T2308 grown in greenhouse (temperature of approximately 25°C, 70 % humidity and natural lighting). Cherries were frozen in liquid nitrogen subsequent to harvesting and stored at -85°C until use. For maturation studies, endosperm and perisperm tissues were separated.

Plant material was ground in liquid nitrogen and extracted in ice cold enzyme extraction buffer (glycerol 10 % v/v, sodium metabisulfite 10 mM, EDTA 5 mM, MOPS (NaOH) 40 mM, pH 6.5) at an approximate ratio of 20 mg per 100 μ l. The mixture was stirred on ice for 20 min, subjected to centrifugation (12,000 g x 30 min), aliquoted and stored at -85°C until use. α -D-galactosidase activity was detected spectrophotometrically with the substrate p-nitrophenyl- α -D-galactopyranoside (pNGP).

The reaction mixture contained 200 μ l pNGP 100 mM in McIlvain's buffer (citric acid 100 mM - Na_2HPO_4 200 mM pH 6.5) up to final volume of 1 ml with enzyme extract. The reaction was maintained at 26°C and started with the addition of enzyme and was stopped by addition of 4 volumes of stop solution (Na_2CO_3 - NaHCO_3 100 mM pH 10.2). Absorption is read at 405 nm. Evolution of nitrophenyl is calculated using molar extinction coefficient $\epsilon = 18300$ (specific for pH 10.2) and converted to $\text{mmol min}^{-1} \text{mg protein}^{-1}$. Total protein was measured in samples extracted in aqueous buffers by the method of Bradford (Anal. Biochem., 72 (1976), 248-254). For the expression of activity, each sample was extracted and aliquoted, and assays were performed in triplicate, the results being expressed as averages.

α -D-galactosidase activity is extremely low or undetected in the young grain stages, and reaches a peak that coincides with a reddening of color of the pericarp. In later stages the activity declines while the cherries are still red. Activities are also compared in different tissues of coffee plant. The activity in perisperm, roots and leaves is particularly low, and close to the limits of detection. However, high activities are recorded in the endosperm where the activity reaches a peak at approximately 36 WAF but also in germinating grain following imbibition of water.

II. Isolation of α -D-galactosidase full-length cDNA from *C. arabica*

Though several coffee α -D-galactosidase cDNA sequences are available in the literature the origin of the coffee material is not indicated. In order see if amino acid and nucleic sequence differences are observed between *C. arabica* and *C. canephora*, it was decided to clone α -D-galactosidase cDNAs from both species.

A cDNA library from *Coffea arabica* var. Caturra T2308 was constructed with polyA⁺ mRNA extracted at 30 weeks after flowering according to Rogers *et al.* (Plant Physiol. Biochem., 37 (1999), 261-272). This plasmid cDNA library (10 ng) was tested by

PCR using the primers BETA1 (SEQ ID NO: 3) and BETA3 (SEQ ID NO: 4) directly deduced from the coffee α -D-galactosidase cDNA sequence (Zhu and Goldstein, 1994). The BETA1 primer is located between the nucleotides 177 and 193 in the sequence SEQ ID NO: 1. The BETA3 primer is located between the nucleotides 1297 and 1313 in the sequence SEQ ID NO: 1. The PCR reaction was performed with *Pfu* DNA polymerase (Stratagene, 11011 North Torrey Pines Road, La Jolla, California 92037, USA) in appropriate 1 X Buffer, 0.2 mM of each dNTP and 0.25 μ M of each oligonucleotides. Denaturation, annealing and extension temperatures are 94 °C for 30 s, 46 °C for 30s and 72 °C for 3 min, respectively. This cycle was repeated 30 times in Robocycler Statagene (USA). PCR products were purified with a Microcon 100 (Millipore SA, BP307 Saint Quentin Yvelines cedex 78054, France) cartridge and ligated in the pCR-Script SK (+) as described by Stratagene (USA). The ligation mixture was then used to transform *E.coli* strain XL1-Blue MRF' and a recombinant vector containing the α -D-galactosidase fragment was purified and cloned into the *Sfr*I site of the pCR-Script SK Amp (+). Its sequence is located between the positions 177 and 1313 of the sequence SEQ ID NO: 1.

According to literature, the 5' end of the α -D-galactosidase cDNA contains 198 bp upstream the 5' end of the BETA 1 primer. In order to clone this sequence from our genotype, we perform an additional PCR, as described previously, with the primers BETA100 (SEQ ID NO: 5) and BETA101 (SEQ ID NO: 6). This *C. arabica* sequence was cloned into the pCR Script Amp SK (+) vector to give the pLP1, and corresponds to SEQ ID NO: 1.

The cDNA obtained contains an open reading frame of 1263 bp, beginning in position 51 and ending in position 1313 of the sequence SEQ ID NO: 1. The translation product corresponds to sequence SEQ ID NO: 2 suggesting that the coffee α -D-galactosidase is synthesized as a preproenzyme, using the translational start codon ATG in position 51 instead of the ATG in position 126. On the other hand, the analysis of the α -D-galactosidase cDNA cloned from *C. canephora* showed that its translation product is very homologous (similarity > 99%) to the protein found in *C. arabica*.

III. Expression of α -D-galactosidase gene during grain maturation

The expression of the gene encoding the α -galactosidase in coffee beans of *C. arabica* Caturra harvested at various stages of development, i.e., 9, 12, 16, 30 and 35 weeks after flowering (WAF) was monitored. To do this, 10 μ g of total RNAs of these coffee beans were

denatured for 15 min at 65°C in 1 x MOPS buffer (20 mM MOPS, 5 mM sodium acetate, 1 mM EDTA, pH 7) in the presence of formamide (50%) and formaldehyde (0.66 M final). They were then separated by electrophoresis, for 6 hrs at 2.5 V/cm, in the presence of 1xMOPS buffer, on a 1.2-% agarose gel containing 2.2 M formaldehyde as final concentration.

After migration, the RNAs were stained with ethidium bromide (BET) according to Sambrook *et al.* (Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press, USA, 1989, chapter 9.31 to 9.51). This makes it possible to standardize the quantities deposited on a gel from the intensities of fluorescence of the 18S and 25S ribosomal RNAs.

The total RNAs were then transferred and fixed on a positively charged Nylon membrane according to the recommendations provided by Boehringer Mannheim (Roche-Boehringer Mannheim GmbH, Biochemica, Postfach 310120, Mannheim 31, DE). The pre-hybridization and hybridization were carried out according to the conditions described above.

Results from Northern-blotting demonstrated a peak of gene expression during the early phase of endosperm development. The peak of specific mRNA expression under greenhouse conditions occurred at approximately 26 WAF, and corresponds to the start of increase in enzyme activity. The peak period of mRNA expression corresponds to the major period of endosperm expansion and hardening taking place in the maturing grains under these conditions. Peak expression for α -galactosidase-specific mRNA either coincided or was slightly later than peak expression of the 11S grain storage protein mRNA. These results led to the construction of an antisense cassette of the coffee cDNA encoding for the α -galactosidase, under the control of the 11S-coffee promoter, in order to reduce the level of α -galactosidase activity in coffee grains under maturation.

IV. Construction of the α -galactosidase antisense cassette

The 11S promoter sequence (Marraccini *et al.*, Plant Physiol. Biochem. 37 (1999), 273-282) from coffee is amplified with the specific primers UP210-1 corresponding to the sequence **SEQ ID NO: 7**, and BAGUS2, corresponding to the sequence **SEQ ID NO: 8**. The oligonucleotide UP210-1 corresponds to the sequence between the nucleotides 24 and 76 published by Marraccini *et al.*, *supra* and contains within its 5' end the synthetic sequence CGGGGTACCCCG containing a *KpnI* restriction site and corresponding to the sequence

SEQ ID NO: 9. The BAGUS2 primer contains in its 5' end the synthetic sequence CGCGGATCCGCG corresponding to the sequence **SEQ ID NO: 10** which carries a *Bam*HI restriction site. This primer also contains the nucleotides 998 to 976 of the sequence published by Marraccini *et al.* (1999). This reaction is carried out in the presence of *Pfu* DNA polymerase (3 units), with 10 ng of pCSPP4 (WO 99/02688), in a final volume of 50 µl containing 10 mM KCl, 6 mM (NH₄)₂SO₄, 20 mM Tris-HCl, pH 8.0, 0.1% Triton X-100, 2 mM MgCl₂, 10 µg/ml BSA, 0.2 mM of each dNTP, 0.25 µM of each oligonucleotides described above. The reaction mixture was then incubated for 30 cycles (94°C-60 s, 55°C-60s, 72°C-3 min) followed by a final extension cycle at 72°C for 7 min.

The PCR fragment of about 950 bp was purified on a Microcon 100 cartridge (Millipore, France), and ligated in the pCR-Script Amp SK (+) vector in the presence of T4 DNA ligase (Promega Corporation, 2800 Woods Hollow Road, Madison, Wisconsin 53711 USA), according to the recommendations provided by the supplier. Next, the *E. coli* strain XL1-Blue MRF' was transformed with the entire ligation mixture. One transformant was selected and its plasmid was purified to sequence the insert in order to determine the orientation of the PCR fragment. This analysis thus made it possible to select the plasmid pLP7.

A shorter version of the 11S promoter was also amplified by the same approach except that the primer UP213-1, having the nucleic sequence **SEQ ID NO: 11** replaces the primer UP210-1. This primer corresponds to the sequence between the nucleotides 754 and 777 published by Marraccini *et al.*, *supra*, and contains in its 5' end the synthetic sequence **SEQ ID NO: 9**. This led to the amplification of a 250 bp fragment of the p11S coffee promoter which was cloned as described previously to give the plasmid pLP8.

The TNOS terminator is amplified following the protocol described for the amplification of the p11S promoter excepted that the primers TNOS1, having the nucleic sequence **SEQ ID NO: 12** and TNOS2, having the nucleic sequence **SEQ ID NO: 13** were used. TNOS1 contains the sequence **SEQ ID NO: 10** in its 5' end. TNOS2 contains the sequence **SEQ ID NO: 9** in its 5' end sequence. These primers led to the amplification of the TNOS sequence from the p35SGFP commercial vector (Clontech Laboratories Inc., 1020 East Meadow Circle, Palo Alto, California 94303-4230 USA). The PCR product was cloned in the pCR-Script Amp SK (+) vector as described before, leading to the recombinant vector called pLP32 and was sequenced to determine its orientation. This vector was then digested by

*Bam*HI to remove the TNOS sequence that was treated afterwards by the T4 DNA polymerase to provide blunt ends.

In the over hand, the pLP7 and pLP8 vectors were linearized by *Eco*RI and also treated with T4 DNA polymerase to provide blunt ends. The TNOS terminator was then cloned in the correct orientation in the pLP7 and pLP8 vectors leading to the vectors p11STNOS7 and p11STNOS7+, respectively.

The α -galactosidase cDNA was amplified from the previously isolated vector pLP1, using the conditions described above except that the primers BETA100B1, having the nucleic sequence **SEQ ID NO: 14** and BETA101B1, having the nucleic sequence **SEQ ID NO: 15** were used. These oligonucleotides correspond to the previously used BETA101 and BETA100 primers in which a *Bam*HI restriction site, corresponding to the sequence **SEQ ID NO: 10**, has been introduced in their 5' ends. The PCR product was cloned into the pCR-Script Amp SK (+) vector as described before, leading to the recombinant vector called pLP20.

This plasmid was digested with *Bam*HI to release the α -galactosidase cDNA. On the other side, recipient vectors p11STNOS7 and p11STNOS7+ were digested independently by means of the same restriction enzyme and dephosphorylated by a CIAP treatment according to the furnisher (Promega, USA). The α -galactosidase cDNA was cloned in the antisense orientation, respectively in the p11STNOS7 and p11STNOS7+ vectors, leading to the vectors designated pALPHA1 and pALPHA9.

To mobilize these cassettes into the binary vector used during the transformation of coffee cell suspension, a final PCR reaction with the *Pfu* DNA polymerase was carried out using the primers UPSAL1, having the nucleic sequence **SEQ ID NO: 16** and UPSAL2, having the nucleic sequence **SEQ ID NO: 17**. Both oligonucleotides contain a *Sal*I restriction site and recognize DNA sequences of the pCR Script Amp SK (+) vector flanking the 11S promoter and the NOS terminator (TNOS) DNA regions. In addition this restriction site is absent from the sequence which was intended to be introduced into the T-DNA of the binary plasmid. These PCR products were cloned again into the pCR-Script Amp SK (+) vector, which was digested with *Sal*I to verify that this restriction site flanks the cassettes. Plasmids obtained were called pALP414 and pALP50, and derive respectively from pALPHA1 and pALPHA9.

V. Cloning of the α -galactosidase antisense cassette in the binary vector of transformation

The α -galactosidase cassettes contained in the vectors pALP414 and pALP50 were sequenced to verify their integrity, particularly to confirm that no point mutations or rearrangements have occurred during the PCR amplification cycles. These cassettes were then purified by digestion of the pALP414 and pALP50 vectors with the *Sa*II restriction enzyme and cloned independently into the pBin19 derivative plasmid related to the vector described by Leroy *et al.* (Plant Cell Rep. **19** (1999), 382-389), except that the gene *cryIAC* was absent. In order to do this, the vector was digested with the *Sa*II restriction enzyme, which recognizes a unique site between the *uidA* and *csrI-1* genes, and was dephosphorylated. After this ligation, the vectors pBIA121, pBIA126 and pBIA9 were selected. In the pBIA121 vector, the *Sa*II cassette obtained from pALP414 is cloned in the orientation [LB] Gus-intron> p11S (long) antisense α -galactosidase cDNA > *csrI-1* [RB]. However, the same cassette is cloned in the reverse orientation in the pBIA126 vector. In the other hand, the *Sa*II cassette obtained from pALP50 cloned in the pBIA9 vector is in the following orientation: [LB] Gus-intron> p11S (short) antisense α -galactosidase cDNA > *csrI-1* [RB].

VI. Transformation of *Agrobacterium tumefaciens*

The binary vectors of transformation pBIA121, pBIA126 and pBIA9 described above were introduced independently into the disarmed *Agrobacterium tumefaciens* strain LBA4404 according to the direct transformation method described by An *et al.* (Plant Mol. Biol. Manuel, Gelvin, Schilperoort and Verma Eds, Kluwer Academic Publishers Dordrecht, Netherlands, **A3** (1993), 1-19). For each transformation, the recombinant *Agrobacterium tumefaciens* clones were selected on LB medium supplemented with kanamycin (50 μ g/ml), streptomycin (100 μ g/ml) and rifampicin (50 μ g/ml).

In order to check the structure of the plasmids introduced into *Agrobacterium tumefaciens*, they were extracted by the rapid mini-preparation technique and were then analyzed by restriction mapping after reverse transformation in *E. coli* strain XL2 Blue MRF⁺.

VII. Transformation of *Coffea* sp.

Leaf explants were cultured and subcultured every five weeks for 3 to 5 months until somatic embryos appeared at the edge of the explants. Somatic embryos were harvested at the torpedo stage, wounded with a sterile scalpel and soaked for two hours in a 0.9% NaCl

solution containing recombinant *Agrobacterium tumefaciens* strain LBA4404 at a OD_{600nm} of 0.3 to 0.5. The co-culture was performed in the dark on semi-solid MS medium without hormones during three days and then washed in liquid MS medium containing cefotaxim (1gr/l) for 3 to 5 hours under constant but gentle agitation. Embryos were cultivated on semi-solid medium with 5µM of BAP, 90µM sucrose in presence of cefotaxim (400mg/l) under low-light condition (16 hrs photoperiod per day). After a period of 3 to 4 weeks, they were transferred to a selective MS medium supplemented with cefotaxim (400mg/l) and chlorsulfuron (80 mg/l). They were then transferred every month to a new selective medium until the regeneration of calli. Transformed embryos growing around the calli were then cultured on the semi-solid MS medium with Morel vitamins (1µM BAP and 30µM sucrose) to induce their germination. After this step, they were transferred on the rooting medium corresponding to the medium described before but without BAP.

To check the effectiveness of the transformation, calli, shoots, roots and leaves were regularly tested for the expression of the *uidA* reporter by a GUS histochemical assay (Jefferson *et al.*, J. EMBO 6 (1987), 3901-3907).

After this procedure, several individual plants were selected and propagated *in vitro* by micro cuttings. Some of them were transferred in greenhouse to achieve their development. No morphological anomalies were observed.

VIII. Analysis of somatic embryos from transformed coffee plants

Somatic embryos were also induced from leaf explants to detect the presence of α-galactosidase antisense mRNA. 11S coffee storage proteins were detected in somatic embryos (Yuffa *et al.*, Plant Cell Rep. 13 (1994) 197-202), suggesting that the *csp1* promoter is active in this tissue. If this is the case, analysis of somatic embryos induced from leaves of young transgenic coffee plants should permit to detect the presence of the α-galactosidase antisense mRNA earlier than in beans.

Total RNAs were then extracted from 100mg of transformed somatic embryos as described previously and tested by RT-PCR using the kit Access RT-PCR system (Promega, USA). Firstly the presence of 11S specific mRNA was confirmed by performing a RT-PCR using the primers located in the coding sequence of the 11S cDNA. This was performed using the primers SO11 corresponding to the sequence **SEQ ID NO: 18** and SO2-1 corresponding to the sequence **SEQ ID NO: 19**. The SO11 primer corresponds to the sequence between the nucleotides 1035 and 1059 of the sequence published by Marraccini *et*

al. , supra. On the other hand, the SO2-1 primer corresponds to the last 24 nucleotides of the sequence published. The synthesis of the first strand of the cDNA (step of reverse transcription) was performed as described by the furnisher (45 min., 48°C). The following parameters were used for the PCR reaction: 45 cycles (60 sec. at 94°C for the denaturation step, 90 sec at 52°C for the annealing step, 4 min at 68°C for the elongation step) with a final extension at 68°C for 7 min.

From this experiment a PCR product of 1590 bp corresponding to the 11S cDNA sequence flanked by the primers SO11 and SO2-1 was obtained. The result confirmed that 11S mRNA were absent from all the tissue tested, i.e. roots, leaves, flowers but were effectively present in somatic embryos of *Coffea canephora* as well as in beans at 27 WAF.

Secondly, the presence of the α -galactosidase sense mRNA was tested by performing a RT-PCR reaction using only the primer B33, corresponding to the sequence **SEQ ID NO: 21** during the phase of reverse transcription (condition 1). This primer corresponds to the complementary sequence of the nucleotides 1286 to 1314 of the sequence **SEQ ID NO: 1**.

In parallel, the detection of the α -galactosidase antisense mRNA was performed using only the primer B11 during the phase of reverse transcription (condition 2). This primer corresponds to the sequence between the nucleotides 50 and 78 of the sequence **SEQ ID NO: 1**. After this reverse transcription (45 min., 48°C), the reaction mixture was treated at 94°C during 1 min to inactivate the MMLV reverse transcriptase. The missing oligonucleotide, B11 in the condition 1 and B33 in condition 2, was added and the reaction was continued by a PCR: 45 cycles (60 sec. at 94°C for the denaturation step, 90 sec at 45°C for the annealing step, 4 min at 68°C for the elongation step) with a final extension at 68°C for 7 min. Using somatic cells from non transformed *C. arabica*, an amplification product of 1310 bp was observed for the condition 1 but not for the condition 2 experiment. However, for somatic embryos obtained from a coffee plantlet transformed by the pBIA9 vector, it was possible to detect an amplification product during the experimental condition 2, confirming the presence of the α -galactosidase antisense mRNA.

Sequence Listings

5

SEQ ID NO: 1

TGCTCCACAAAGCAGTGGCAATTGAGTTGATTGATCAACACCAATTTACCATGGCCGCTGCTT
ATTACTACCTTTTTTCTAGTAAAAAAGCCACCAAAGCTGGTGCTCCGAGCTTCGTTATTGA
TGTTTTTATGTTTCTTGGCGGTTGAAAACGTTGGTGCTTCCGCTCGCCGGATGGTGAAGTCTC
10 CAGGAACAGAGGATTACACTCGCAGGAGCCTTTTAGCAAATGGGCTTGGTCTAACACCACCGA
TGGGGTGGAAACAGCTGGAATCATTTAGTTGTAATCTTGATGAGAAATTGATCAGGGAAACAG
CCGATGCAATGGCATCAAAGGGGCTTGCTGCACTGGGATATAAGTACATCAATCTTGATGACT
GTTGGGCAGAACTTAACAGAGATTCACAGGGGAATTTGGTTCTTAAAGGTTCAACATTCCCAT
CAGGGATCAAAGCCTTAGCAGATTATGTTACAGCAAAGGCCTAAAGCTTGGAATTTACTCTG
15 ATGCTGGAACCTCAGACATGTAGTAAACTATGCCAGGTTCAATAGGACACGAAGAACAAGATG
CCAAAACCTTTGCTTCATGGGGGGTTGATTACTTAAAGTATGACAACTGTAACGACAACAACA
TAAGCCCCAAGGAAAGGTATCCAATCATGAGTAAAGCATTGTTGAACTCTGGAAGGTCCATAT
TTTTCTCTCTATGTGAATGGGGAGATGAAGATCCAGCAACATGGGCAAAGAAGTTGGAAACA
GTTGGAGAACCACTGGAGATATAGATGACAGTTGGAGTAGCATGACTTCTCGGGCAGATATGA
20 ACGACAAATGGGCATCTTATGCTGGTCCCGGTGGATGGAATGATCCTGACATGTTGGAGGTGG
GAAATGGAGGCATGACTACAACGGAATATCGATCCCATTTAGCATTTGGGCATTAGCAAAG
CACCTCTACTGATTGGCTGTGACATTCGATCCATTGACGGTGCGACTTTCCAACCTGTTAAGCA
ATGCGGAAGTTATTGCGGTTAACCAAGATAAACTTGCGGTTCAAGGGAAAAGGTTAAGACTT
ACGGAGATTTGGAGGTGTGGGCTGGACCTCTTAGTGGAAGAGAGTAGCTGTGCTTTGTGGA
25 ATAGAGGATCTTCCACGGCTACTATTACCGCGTATTTGGTCCGACGTAGGCCTCCCGTCCACGG
CAGTGGTTAATGCACGAGACTTATGGGCGCATTCAACCGAAAAATCAGTCAAAGGACAAATCT
CAGCTGCAGTAGATGCCACGATTGAAAATGTATGTCCTAACCCACAGTGATTAACAGGAG
AATGCAGAAGACAAGTGATGGTTGGCTCTTTCAAGGATTTGATTACCTTAAAGAATTTTTTAC
ATGTTATGAATCAATTCAAAGCAATTATGTGTTTTGAAGAGATTAAGTCAATAAAT

30

SEQ ID NO: 2

MAAAYYYLFSSKKSHQKLVLRASLLMFLCFLAVENVGASARRMVKSPGTEDYTRRS
LLANGLGLTPPMGWNSWNHFCNLDEKLIRETADAMASKGLAALGYKYINLDDCW
AELNRDSQGNLVPKGSTFPSGIKALADYVHSGKGLKLGISDAGTQTCSKTMPSGLGH
35 EEQDAKTFASWGVLDYLKYDNCNDNNISPKERYPIMSKALLNSGRSIFSLCEWGED

PATWAKEVGNSWRTTGDIDDSWSSMTSRADMNDKWASYAGPGGWNDPDMLEV
NGGMTTTEYRSHFSIWALAKAPLLIGCDIRSIDGATFQLLSNAEVIAVNQDKLGVQG
KKVKTYGDLEWAGPLSGKRVAVALWNRGSSTATITAYWSDVGLPSTAVVNARDL
WAHSTEKSVKGQISAAVDAHDSKMYVLTPQ

5

SEQ ID NO: 3 (BETA1)

ATGGTGAAGTCTCCAGG

SEQ ID NO: 4 (BETA3)

10 TCACTGTGGGGTTAGGA

SEQ ID NO: 5 (BETA 100)

TGCTCCACAAAGCAGTGGCAATT

15 SEQ ID NO: 6 (BETA101)

ATTTATTGACTTAATCTCTTCAA

SEQ ID NO 7: 210-1

CGGGGTACCCCGCCTCTTTTCTTTTGGAGTACAAG

20

SEQ ID NO 8: BAGUS 2

CGCGGATCCGCGTCTCTGACAACAGAGGAGAGTGT

SEQ ID NO: 9. *KpnI*

25 CGGGGTACCCCG

SEQ ID NO: 10. *BamHI*

CGCGGATCCGCG

30 SEQ ID NO 11: UP213-1

CGGGGTACCCCGACAAAAGATTGAACAATACATGTC

SEQ ID NO 12: TNOS1

CGCGGATCCGCGGAGCTCGAATTCCCCGATCGTT

5 SEQ ID NO 13: TNOS2

CGGGGTACCCCGGAATTCCCGATCTAGTAACATAG

SEQ ID NO 14: BETA100BI

CGCGGATCCGCGTGCTCCACAAAGCAGTGGCAATT

10

SEQ ID NO 15: BETA101BI

CGCGGATCCGCGATTATTGACTTAATCTCTTCAA

SEQ ID NO 16: UPSAL1

15 CACGCGTCGACGCTCCACCGCGGTGGCGGCCGCTC

SEQ ID NO 17: UPSAL2

GGGCCCCCCTCGAGGTCGACGG

20 SEQ ID NO 18: SO11

TTCITTTGTTCCTCGGCTGTTTG

SEQ ID NO 19: SO2-1

CCAAACATCAAACCTTCTCGCAATC

25

SEQ ID NO: 20 (BETA11)

ATGGCCGCTGCTTATTACTACCTTTTT

SEQ ID NO: 21 (BETA33)

30 TCACTGTGGGGTTAGGACATACATTTT

SEQUENCE LISTING

<110> Société des Produits Nestlé S.A.
 <120> Coffee plant with reduced alpha-D-galactosidase activity
 <130> 80331WO
 <150> EP 01124160
 <151> 2001-10-10
 <160> 21
 <170> PatentIn version 3.1
 <210> 1
 <211> 1442
 <212> DNA
 <213> Coffea arabica
 <400> 1
 tgctccacaa agcagtggca attgagttga ttgatcaaca ccaatttacc atggccgctg 60
 cttattacta ccttttttct agtaaaaaaa gccacaaaaa gctggtgctc cgagcttcgt 120
 tattgatgtt tttatgtttc ttggcggttg aaaacgttgg tgcttccgct cgccggatgg 180
 tgaagtctcc aggaacagag gattacactc gcaggagcct ttagcaaat gggcttggtc 240
 taacaccacc gatggggtgg aacagctgga atcatttcag ttgtaatctt gatgagaaat 300
 tgatcaggga aacagccgat gcaatggcat caaaggggct tgctgcactg ggatataagt 360
 acatcaatct tgatgactgt tgggcagaac ttaacagaga ttcacagggg aatttggttc 420
 ctaaagggtc aacattccca tcagggatca aagccttagc agattatgtt cacagcaaat 480
 gcctaaagct tggaatttac tctgatgctg gaactcagac atgtagtaaa actatgccag 540
 gttcattagg acacgaagaa caagatgccaa aaacctttgc ttcattggggg gttgattact 600
 taaagtatga caactgtaac gacaacaaca taagcccaa ggaaagggtat ccaatcatga 660
 gtaaagcatt gttgaactct ggaagggtcca tatttttctc tctatgtgaa tggggagatg 720
 aagatccagc aacatgggca aaagaagttg gaaacagttg gagaaccact ggagatatag 780
 atgacagttg gagtagcatg acttctcggg cagatatgaa cgacaaatgg gcatcttatg 840
 ctgggtcccg tggtatggaat gatcctgaca tggtggaggt gggaaatgga ggcatgacta 900
 caacggaata tcgatcccat ttcagcattt gggcattagc aaaagcacct ctactgattg 960
 gctgtgacat tcgatccatt gacggtgcca ctttccaact gttaagcaat gcggaagtta 1020
 ttgcggttaa ccaagataaa cttggcggtc aagggaaaaa ggttaagact tacggagatt 1080
 tggaggtgtg ggctggacct cttagtggaa agagagtagc tgctgctttg tggaatagag 1140
 gatcttccac ggctactatt accgcgtatt ggtccgacgt aggcctcccg tccacggcag 1200
 tgggtaatgc acgagactta tgggcgcatt caaccgaaaa atcagtcaaa ggacaaatct 1260

cagctgcagt agatgccac gattcgaaaa tgtatgtcct aacccacag tgattaacag 1320
 gagaatgcag aagacaagtg atggttggct ctttcaagga tttgattacc ttaaagaatt 1380
 tttcacatgt tatgaatcaa ttcaaagcaa ttatgtgttt tgaagagatt aagtcaataa 1440
 at 1442

<210> 2
 <211> 420
 <212> PRT
 <213> Coffea arabica

<400> 2

Met Ala Ala Ala Tyr Tyr Tyr Leu Phe Ser Ser Lys Lys Ser His Gln
 1 5 10 15

Lys Leu Val Leu Arg Ala Ser Leu Leu Met Phe Leu Cys Phe Leu Ala
 20 25 30

Val Glu Asn Val Gly Ala Ser Ala Arg Arg Met Val Lys Ser Pro Gly
 35 40 45

Thr Glu Asp Tyr Thr Arg Arg Ser Leu Leu Ala Asn Gly Leu Gly Leu
 50 55 60

Thr Pro Pro Met Gly Trp Asn Ser Trp Asn His Phe Ser Cys Asn Leu
 65 70 75 80

Asp Glu Lys Leu Ile Arg Glu Thr Ala Asp Ala Met Ala Ser Lys Gly
 85 90 95

Leu Ala Ala Leu Gly Tyr Lys Tyr Ile Asn Leu Asp Asp Cys Trp Ala
 100 105 110

Glu Leu Asn Arg Asp Ser Gln Gly Asn Leu Val Pro Lys Gly Ser Thr
 115 120 125

Phe Pro Ser Gly Ile Lys Ala Leu Ala Asp Tyr Val His Ser Lys Gly
 130 135 140

Leu Lys Leu Gly Ile Tyr Ser Asp Ala Gly Thr Gln Thr Cys Ser Lys
 145 150 155 160

Thr Met Pro Gly Ser Leu Gly His Glu Glu Gln Asp Ala Lys Thr Phe
 165 170 175

Ala Ser Trp Gly Val Asp Tyr Leu Lys Tyr Asp Asn Cys Asn Asp Asn
 180 185 190

Asn Ile Ser Pro Lys Glu Arg Tyr Pro Ile Met Ser Lys Ala Leu Leu
 195 200 205
 Asn Ser Gly Arg Ser Ile Phe Phe Ser Leu Cys Glu Trp Gly Asp Glu
 210 215 220
 Asp Pro Ala Thr Trp Ala Lys Glu Val Gly Asn Ser Trp Arg Thr Thr
 225 230 235 240
 Gly Asp Ile Asp Asp Ser Trp Ser Ser Met Thr Ser Arg Ala Asp Met
 245 250 255
 Asn Asp Lys Trp Ala Ser Tyr Ala Gly Pro Gly Gly Trp Asn Asp Pro
 260 265 270
 Asp Met Leu Glu Val Gly Asn Gly Gly Met Thr Thr Thr Glu Tyr Arg
 275 280 285
 Ser His Phe Ser Ile Trp Ala Leu Ala Lys Ala Pro Leu Leu Ile Gly
 290 295 300
 Cys Asp Ile Arg Ser Ile Asp Gly Ala Thr Phe Gln Leu Leu Ser Asn
 305 310 315 320
 Ala Glu Val Ile Ala Val Asn Gln Asp Lys Leu Gly Val Gln Gly Lys
 325 330 335
 Lys Val Lys Thr Tyr Gly Asp Leu Glu Val Trp Ala Gly Pro Leu Ser
 340 345 350
 Gly Lys Arg Val Ala Val Ala Leu Trp Asn Arg Gly Ser Ser Thr Ala
 355 360 365
 Thr Ile Thr Ala Tyr Trp Ser Asp Val Gly Leu Pro Ser Thr Ala Val
 370 375 380
 Val Asn Ala Arg Asp Leu Trp Ala His Ser Thr Glu Lys Ser Val Lys
 385 390 395 400
 Gly Gln Ile Ser Ala Ala Val Asp Ala His Asp Ser Lys Met Tyr Val
 405 410 415
 Leu Thr Pro Gln
 420

<210> 3
 <211> 17

<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA1

<400> 3
atggtgaagt ctccagg

17

<210> 4
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA3

<400> 4
tcactgtggg gtttaga

17

<210> 5
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA100

<400> 5
tgctccacaa agcagtggca att

23

<210> 6
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA101

<400> 6 --
atttattgac ttaatctctt caa

23

<210> 7
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer 210-1

<400> 7
cggggtaccc cgcctctttt cttttggagt acaag

35

<210> 8
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
 <223> Primer BAGUS2

 <400> 8
 cgcggtatccg cgtctctgac aacagaggag agtgt

35

<210> 9
 <211> 12
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> KpnI

 <400> 9
 cggggtaccc cg

12

<210> 10
 <211> 12
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> BamHI

 <400> 10
 cgcggtatccg cg

12

<210> 11
 <211> 36
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Primer UP213-1

 <400> 11
 cggggtaccc cgacaaaaga ttgaacaata catgtc

36

<210> 12
 <211> 35
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Primer TNOS1

 <400> 12
 cgcggtatccg cggagctcga atttccccga tcgtt

35

<210> 13
 <211> 35
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Primer TNOS2

<400> 13
 cggggtaccc cggaattccc gatctagtaa catag 35

<210> 14
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer BETA100BI

<400> 14
 cgcggatccg cgtgctccac aaagcagtgg caatt 35

<210> 15
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer BETA101BI

<400> 15
 cgcggatccg cgatttattg acttaatctc ttcaa 35

<210> 16
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer UPSAL1

<400> 16
 cacgcgtcga cgctccaccg cggcggcggc cgctc 35

<210> 17
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer UPSAL2

<400> 17
 gggccccccc tcgaggtcga cgg 23

<210> 18
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer S011

<400> 18
 ttcttttgtt cctcggctgt ttg 23

<210> 19
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer SO2-1

<400> 19
ccaacatca aacttctcgc aatc

24

<210> 20
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA11

<400> 20
atggcgcgtg cttattacta ccttttt

27

<210> 21
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer BETA33

<400> 21
tcactgtggg gttaggacat acatttt

27